**References / Bibliography**


**1.** World Health Organization. (1969). WHO Expert Committee on Drug Dependence. Sixteenth report. (Technical report series. No. 407).Geneva: World Health Organization.

**2.** Rother K (April 2007). "Diabetes treatment—bridging the divide". The New England Journal of Medicine 356 (15): 1499–501.

**3.** Shoback, edited by David G. Gardner, Dolores (2011). Greenspan's basic & clinical endocrinology (9th ed. ed.). New York: McGraw-Hill Medical. pp. Chapter 17.

**4.** Cooke DW, Plotnick L (November 2008). "Type 1 diabetes mellitus in pediatrics". Pediatr Rev 29 (11): 374–84).

**5.** K.S Lakshmi, T. Rajesh, Shrinivas Sharma, "Simultaneous determination of metformin and pioglitazone byreversed phase HPLC in pharmaceutical dosage forms". International Journal of Pharmacy and Pharmaceutical Sciences, Vol. 1, Issue 2, Oct-Dec. 2009.

**6.** Adukondalu D, P S Malathy, J Venkateshwar rao, Y Madhusudan Rao. Development and Validation of HPLC method for detection of pioglitazone hydrochloride in dosage forms. IJPBS,Volume 1, Issue 4, oct-dec 2011. 474-478.

**7.** Zarghi A, Foroutan SM, Shafaati A, Khoddam A. Rapid determination of metformin in human plasma using ion-pair HPLC. J Pharm Biomed Anal. 2003 Feb 5;31(1):197-200.

**8.** Vasudevan M, Ravi J, Ravisankar S, Suresh B. Ion-pair liquid chromatography technique for the estimation of metformin in its multicomponent dosage forms.J Pharm Biomed Anal. 2001 Apr;25(1):77-84.

**9.** Sahoo P. K., R. Sharma, and S. C. Chaturvedi Simultaneous Estimation of Metformin Hydrochloride and Pioglitazone Hydrochloride by RP-HPLC Method from Combined Tablet Dosage Form Indian J Pharm Sci. 2008 May-Jun; 70(3): 383–386.

**10.** Xiang FeiJiu, De Wei Shang, Ye Chen, XinGangLi, Xiao Meng Wan, Tian Yan Zhou, Wei Lu. A high performance liquid chromatography method for the quantitative determination assay of sitagliptin in rat plasma and its applicationin pharmacokinetics study X. F. Jiu et al. / Journal of Chinese Pharmaceutical 64 Sciences 20 (2011) 63–69.

**11.** Cristina georgiță1, Iulia sora1, Florin albu, Crina maria monciu. Comparison of a LC/MS method with a LC/UV method for the determination of metformin in plasma samples. FARMACIA, 2010, Vol.58, 2.

**12.** Changkang Pan, Frances Liu, Michael Motto. Identification of pharmaceutical impurities in formulated dosage forms. Journal of Pharmaceutical Sciences, Volume 100, Issue 4, pages 1228–1259, April 2011.

**13.** Rathinavel G., U.umanath, J.valarmathy, L.samueljoshua, C.selvin thanuja, M.ganesh, T.sivakumar and R.priyadarsini. RP-HPLC Method for the Simultaneous Estimation of Rosiglitazone and Gliclazide in Tablets. E-Journal of Chemistry 2009, 6(4), 1188-1192.

**14.** Bhamare P.C., S.B.Bari, S.Natarajan, A.A.Patil, S.H.Patil, P.T.Shirode. Development and Validation of a precise single stability indicating HPLC method for determinations of Metformin hydrochloride and Fenofibrate, in Pure form and in Pharmaceutical tablets. International Journal of Pharm Tech Research, Vol. 3, No.1, pp 505-515, Jan-Mar 2011.

**15.** Anand Prem D.C., K.L.Senthil kumar, B.Senthil kumar, M.Saravana kumar, R.Thirumurthy. A New RP-HPLC Method Development and Validation for Simultaneous estimation of Telmisartan and Pioglitazone in Pharmaceutical Dosage Form. International Journal of Chem Tech Research, Vol. 3, No.1, pp 448-454, Jan-Mar 2011.

**16.** Wei Zeng, Donald G. Musson, Alison L. Fisher, Li Chen,Michael S. Schwartz,Eric J. Woolf,Amy Qiu Wang. Determination of sitagliptin in human urine and hemodialysate using turbulent flow online extraction and tandem mass spectrometry. Journal of Pharmaceutical and Biomedical Analysis Volume 46, Issue 3, 13 February 2008, Pages 534–542.

**17.** Malleswararao Chellu S. N., Mulukutla V. Suryanarayana, Khagga mukkanti. Simultaneous Determination of Sitagliptin Phosphate Monohydrate and Metformin Hydrochloride in Tablets by a Validated UPLC Method. Sci Pharm. 2012; 80: 139–152.

**18.** Ramzia I. El-Bagary, Ehab F. Elkady, Bassam M. Ayoub. Liquid Chromatographic Methods for the Determination of Vildagliptin in the Presence of its Synthetic Intermediate and the Simultaneous Determination of Pioglitazone Hydrochloride and Metformin Hydrochloride. International Journal of Biomedical Science 7(3), 201-208, Sep 15, 201.

**19.** Rashmitha N., Hiriyanna S.G , CH. Sreenivasa Rao , K. Chandra Sekhar Reddy , M. Hari Kiran , Hemant Kumar Sharmaa , K. Mukkanti, A validated stability indicating HPLC method for the determination of impurities in pioglitazone hydrochloride. Der Pharma Chemica, 2010, 2(5): 426-433.

**20.** AMR Lotfy Saber, Determination of Pioglitazone Hydrochloride in Tablets by High-Performance Liquid Chromatography, Pak. J. Anal. Environ. Chem. Vol. 9, No. 2 (2008) 118 . 121.

**21.** Srinivasulu D., B.S.sastry, G.Omprakash, Development and Validation of new RP-HPLC method for determination of pioglitazone HCl in pharmaceutical dosage forms Int J Chem Res, Vol 1, issue 1 (2010), 1820.

**22.** E. Fedorov; J.-F. Larocque; A. Pimenov; M. Mancini Fast LC-MS-MS Determination of Metformin in Human Plasma with Simultaneous Liquid-Liquid Extraction and Derivatization. WARNEX Bioanalytical Services, Laval, QC, Canada.

**23.** Patil Mahendra K., Ajay S. Raghuwanshi1, U. K. Jain, Ajit S. Raghuwanshi, Nilesh Gupta and Ajay Patel, RP-HPLC Method for the Determination of Rosiglitazone in presence of its Degradation Products in Bulk Drugs, Der Pharmacia Sinica, 2011, 2 (2):368-374.

**24.** Patrícia Gomes, Juliana Sippel, André Jablonski, Martin Steppe. Determination of rosiglitazone in coated tablets by MEKC and HPLC method. Journal of Pharmaceutical and Biomedical Analysis, Volume 36, Issue 4, 19 November 2004, Pages 909–913.

**25.** Ravikanth CH**,** A. Anil Kumar, V Uday Kiran, S Prashanth, B Madhu, Y Narsimha Reddy. Sensitive and Rapid HPLC Method for the Determination of Pioglitazone in Rat Serum. International Journal of Pharmaceutical Sciences and Drug Research 2011; 3(1): 38-41.

**26.** Jedlicka A, Klimes J, Grafnetterová T. Reversed-phase HPLC methods for purity test and assay of pioglitazone hydrochloride in tablets. Pharmazie. 2004 Mar;59 (3):178-82.

**27.** Pattana Sripalakit, Penporn Neamhom, Aurasorn Saraphanchotiwitthaya High-performance liquid chromatographic method for the determination of pioglitazone in human plasma using ultraviolet detection and its application to a pharmacokinetic study. Journal of Chromatography B. Volume 843, Issue 2, 7 November 2006, Pages 164–169.

**28.** Rashmitha N., Hiriyanna S.G, CH. Sreenivasa Rao, K. Chandra Sekhar Reddy, M. Hari Kiran, Hemant Kumar Sharma, K. Mukkanti. A Validated Stability Indicating HPLC method for the Determination of Impurities in Pioglitazone Hydrochloride**.** Der Pharma Chemica, 2010, 2(5): 426-433.

**29.** Gebremriam Ketema, D Gowris Sankar, "Development and validation of rp-HPLC method for simultaneous estimation of sitagliptin and simvastatin in

bulk and tablet dosage forms", Inventi Impact: Pharm Ana & Qual Assur, Vol. 2012 ,278/12, 2012.

**30.** Shyamala.M, Mohideen.S, Satyanarayana .T, Ch. Narasimha Raju, Suresh Kumar.P, Swetha.K.Validated RP-HPLC for simultaneous estimation of sitagliptin phosphate and metformin hydrochloride in tablet dosage form. AJPTR. 2011; 1(2): 93-101.

**31.** Stella H. Vincent, James R. Reed, Arthur J. Bergman, Charles S. Elmore, Bing Zhu, Shiyao Xu, David Ebel, Patrick Larson, Wei Zeng, Li Chen, Stacy Dilzer, Kenneth Lasseter, Keith Gottesdiener, John A. Wagner and Gary A. Herman, Metabolism And Excretion of the Dipeptidyl Peptidase 4 Inhibitor [$^{14}$C] Sitagliptin in Humans. drug metabolism and disposition vol. 35, no. 4.